

Growth Potential of *Bacillus cereus* Group Strains from Different Phylogenetic Groups in a Dairy Food Model

Tyler Chandross-Cohen¹, Mackenna Yount¹, Jun Su², Chenhao Qian², Martin Wiedmann², Jasna Kovac¹

¹Department of Food Science, The Pennsylvania State University, University Park, PA 16802

²Department of Food Science, Cornell University, Ithaca, NY 14850



Background and Objectives



The *Bacillus cereus* group are Gram-Positive, closely related spore-forming bacteria. Some members of this group are cytotoxic and cause foodborne illness, while others are used as bioinsecticides or probiotics in the agricultural industry¹.



The range of growth temperatures of the *B. cereus* group varies between *panC* phylogenetic clades².



Understanding the differences in the growth capabilities of cytotoxic *B. cereus* group strains belonging to different phylogenetic clades is needed to inform the development of exposure and risk assessment models in milk³.

Objectives:

- (1) Evaluate the growth in skim milk broth (SMB) of spores from 17 cytotoxic *B. cereus* group isolates with unique virulence gene profiles and *panC* phylogenetic clades.
- (2) Compare growth data from room temperature (22°C), exceeded refrigeration temperature (10°C), and refrigeration temperature (4°C) to evaluate differences between clades in their maximum population size, doubling time, and the maximum growth rate.

Materials and Methods




Figure 1: Isolates were streaked onto brain heart infusion agar (BHI), incubated, and inoculated into BHI broth. Sporulation was induced using AK #2 agar for 5 days. Phase contrast microscopy was used to confirm the presence of spores. Spores were harvested and suspended in sterile deionized water. Suspensions were quantified in CFU/ml.




Figure 2: Spore suspensions were heat-treated at 80°C to promote germination and kill remaining vegetative cells. Spore suspensions were inoculated into sterile SMB at $\sim 10^3$ spores/ml. The skim milk broth was sampled at the set time points based on the temperature. Dilutions were spiral plated onto BHI and counted using the Neutec SphereFlash colony scanner.

Conclusions

- (1) At both 22°C and 10°C, the growth of *B. cereus* species exhibited significant differences among phylogenetic groups.
- (2) Tested *B. cereus* isolates did not grow at 4°C.
- (3) Most isolates in this study grew above 10^5 CFU/ml at exceeded refrigeration temperature (10°C), which has previously been associated with foodborne disease⁴.

Significance

Accurately predicting the growth of *B. cereus* species in milk is imperative to help mitigate the risk of foodborne illnesses and protect public health

Future Directions

- (1) Use growth data to build an exposure assessment model for milk
- (2) Test more isolates representing individual phylogenetic groups to improve the accuracy of the exposure assessment model
- (3) Evaluate growth in another food model

References

- (1) Carroll, L. M. et al. (2022). *Critical Reviews in Food Science and Nutrition*, 62 (28), 7677–7702.
- (2) Ceuppens, S. et al. (2013). *FEMS Microbiology Ecology*, 54 (3), 433–450.
- (3) Martin, N. H. et al. (2012). *J Dairy Sci* 2012, 95 (12), 7384–7390.
- (4) Tewari, A. et al. (2015). *J Food Sci Technol*, 52 (5), 2500–2511.

Results




Figure 3: Growth of *B. cereus* species from different clades (I, II, III, IV, V, VII) in SMB at 22°C, 10°C, and 4°C. The Growthcurver package in R was used to collect growth metrics. There was a significant difference ($\alpha=0.05$) in the estimated maximum population size ($p=0.0058$), doubling time ($p<0.001$), and the maximum growth rate ($p<0.001$) between isolates grown at 10°C compared to 22°C. Isolates from *panC* clade I grew significantly less ($\alpha=0.05$) at 22°C ($p=0.00139$) and 10°C ($p=7.76 \times 10^{-7}$) compared to all other clades.